

## RESEARCH ARTICLE

# Sleep duration and physical performance during a 6-week military training course

David T. Edgar<sup>1,2</sup>  | Nicholas D. Gill<sup>1</sup> | Christopher Martyn Beaven<sup>1</sup>  |  
Jennifer L. Zaslona<sup>4</sup>  | Matthew W. Driller<sup>3</sup> 

<sup>1</sup>Division of Health, Engineering, Computing & Science, Te Huataki Waiora School of Health, University of Waikato, Tauranga, New Zealand

<sup>2</sup>New Zealand Defence Force, Wellington, New Zealand

<sup>3</sup>Sport and Exercise Science, School of Allied Health, Human Services and Sport, La Trobe University, Melbourne, Vic., Australia

<sup>4</sup>Sleep/Wake Research Centre, Massey University, Wellington, New Zealand

## Correspondence

David T. Edgar, Division of Health, Engineering, Computing & Science, Te Huataki Waiora School of Health, University of Waikato, Tauranga, New Zealand.

Email: David.Edgar@nzdf.mil.nz

## Summary

Sleep is vital in influencing effective training adaptations in the military. This study aimed to assess the relationship between sleep and changes in physical performance over 6 weeks of military training. A total of 22 officer-trainees (age:  $24 \pm 5$  years) from the New Zealand Defence Force were used for this prospective cohort study. Participants wore wrist-actigraphs to monitor sleep, completed subjective wellbeing questionnaires weekly, and were tested for: 2.4-km run time-trial, maximum press-up and curl-ups before and after 6 weeks of training. Average sleep duration was calculated over 36 nights ( $6:10 \pm 0:28$  hr:min), and sleep duration at the mid-point ( $6:15$  hr:min) was used to stratify the trainees into two quantile groups (UNDERS:  $5:51 \pm 0:29$  hr:min,  $n = 11$ ) and (OVERS:  $6:27 \pm 0:09$  hr:min,  $n = 11$ ). There were no significant group  $\times$  time interactions for 2.4-km run, press-ups or curl-ups ( $p > .05$ ); however, *small* effects were observed in favour of OVERS for 2.4-km run (59.8 versus 44.9 s;  $d = 0.26$ ) and press-ups (4.7 versus 3.2 reps;  $d = 0.45$ ). Subjective wellbeing scores resulted in a significant group  $\times$  time interaction ( $p < .05$ ), with *large* effect sizes in favour of the OVERS group for Fatigue in Week 1 ( $d = 0.90$ ) and Week 3 ( $d = 0.87$ ), and Soreness in Week 3 ( $d = 1.09$ ) and Week 4 ( $d = 0.95$ ). Sleeping more than  $6:15$  hr:min per night over 6 weeks was associated with *small* benefits to aspects of physical performance, and *moderate* to *large* benefits on subjective wellbeing measures when compared with sleeping  $< 6:15$  hr:min.

## KEYWORDS

actigraphy, armed forces, exercise recovery, sleep restriction

## 1 | INTRODUCTION

An estimated 75% of young adults sleep  $< 8$  hr per night (Owens, 2014). Sleep deficiency is of growing concern for the general public (Brown et al., 2013), and for the military with the recruitment of young adults (Good et al., 2020; Miller & Shattuck, 2005). Military personnel can experience even greater challenges with sleep due to the stressful and constantly changing nature of daily training and operation (Good et al., 2020). Sleep

management in the military can be complicated due to the need to undertake tasks both day and night at very short notice (Williams et al., 2014). In a consensus paper by Lovalekar et al. (2018), sleep was identified as an emerging research priority area at the International Congress on Soldiers Physical Performance, ranking third out of 43 topics identified by 502 attendees from 32 countries at the congress.

It has been established that sleep can be negatively affected during military training (Williams et al., 2014), and further impacts

physical performance when deployed (Brown et al., 2013), especially when below the recommendation of 7–9 hr per night for adults stated in the joint consensus of the American Academy of Sleep Medicine and Sleep Research Society (Watson et al., 2015). A review paper by Miller et al. (2012) discussed the effects of sleep deprivation on human performance, and outlined research showing that short sleep duration has a negative effect on operational physical performance tasks in the military, such as: carrying and lifting, patrolling over distance, weapons handling, and equipment control. It has also been suggested that a lack of sleep may contribute to reduced gains in physical performance and increased injury occurrence in the military (Lentino et al., 2013; Miller et al., 2012). Consecutive days of reactive operation can also diminish task effectiveness due to an adverse effect on sleep quality (Miller et al., 2011; Williams et al., 2014). Chronically sleeping less than the recommended 7–9 hr per night has also been reported to negatively impact physical performance, and can contribute to fatalities during military operations (Williams et al., 2014).

Therefore, the importance of sleep and its role in recovery and enhancing physical performance in the military is of the utmost importance (Brown et al., 2013). Williams et al. (2014) determined that insufficient sleep occurs during both basic training and during academic phases of study at military academies, and identified that United States (US) Military Academy cadets had an average weekday sleep duration of <5 hr per night, and average weekend sleep duration of ~6.5 hr (Williams et al., 2014). Previous research supports how the recommended duration of sleep generally does not occur in the military environment (Good et al., 2020; Lentino et al., 2013; Moore et al., 2020). The need for further research on the long-term effect of lack of sleep on adaptation to training and physical development in military personnel has also been highlighted (Miller et al. 2010). Lentino et al. (2013) found that short sleepers were less likely to have a healthy body composition, meet physical training (PT) recommendations, and pass their Army physical fitness tests.

The current body of literature is limited when assessing the effect of sleep on long-term physical performance adaptation in the military. There is also limited information on the use of objective sleep measures (e.g. wrist-actigraphy) in military settings. Therefore, the purpose of this study was to investigate the relationship between sleep (via wrist-actigraphy), physical performance, and the subjective wellbeing of officer trainees during 6 weeks of initial military training.

## 2 | METHODS

### 2.1 | Participants

A total of 22 healthy officer trainees, a representative sample of the New Zealand Defence Force (NZDF) Joint Officer Induction Course (JOIC), participated in the current study (Table 1). Participation in the study was voluntary with inclusion dependent on passing the pre-course medical examination. Trainee's data were to be excluded if they withdrew voluntarily from the course, or were medically removed. Ethical approval for the study was obtained from the University of Waikato Human Research Ethics Committee (HREC) (Health) #2018-01.

### 2.2 | Study design

The study design was a prospective cohort study, whereby sleep was monitored via wrist-actigraphy in all trainees over a 6-week JOIC, with physical performance assessed pre- and post-training. Fitness and performance data were collected in Weeks 1 and 6 of the JOIC across two 90-min sessions, with subjective wellness questionnaires collected weekly. The sleep of trainees was monitored via wrist-actigraphy for the entire duration of the 6-week course in order to assess the relationship between sleep and changes in physical performance.

### 2.3 | Wrist-actigraphy

Wrist-actigraphy has previously been used as a practical sleep assessment method in a military environment (Kushida et al., 2001). A Micro Motionlogger<sup>®</sup> (Ambulatory Monitoring) Sleep Watch was allocated to every trainee, data were collected using the device's zero-crossing mode and recorded in 1-min epochs, with individual devices worn continuously for the full duration of the course during both wake and sleep on whichever wrist felt comfortable (Driller et al., 2017). The validity and reliability of the Micro Motionlogger has previously been reported and deemed acceptable (Tryon, 2008). Sleep and wakefulness were inferred based on activity count using the Cole-Kripke software algorithm for sleep estimation using the methods described by Quante (2018). This technique using the AMI software analysis has previously been

TABLE 1 Participant demographics for the included and excluded sample

Group	No.	Male	Female	Age	Weight (kg)	Height (cm)
Sample	22	19	3	24 ± 5	81 ± 28	180 ± 18
Non-participants	72	55	17	22 ± 5	78 ± 18	177 ± 8
Full cohort	94	74	20	23 ± 4	79 ± 12	178 ± 9

Data shown as mean ± SD.

compared with polysomnography, and shown to correctly distinguish sleep from wakefulness approximately 88% of the time (Morgenthaler, 2007). The device was removed for any water submersion activities and placed back on the wrist immediately post-activity. Double scoring by two trained members of the research team was undertaken on 33% of randomly selected sleep files to assess the reliability of manual selection of sleep intervals. Any discrepancies of more than 15 min for either “start time” or “end time” of the sleep interval were flagged and re-analysed. A good interrater reliability agreement rate of 88% was achieved (McHugh, 2012).

## 2.4 | Subjective wellbeing monitoring

During the study period, trainees completed a psychological wellbeing questionnaire at the end of each week that was based on the recommendations of Hooper and Mackinnon (1995). The questionnaire assessed each trainee's fatigue, sleep quality, general muscle soreness, stress levels and mood on a five-point scale of 1–5 with 0.5-point increments (5 = very good, 4 = good, 3 = normal, 2 = poor, 1 = very poor).

## 2.5 | PT programme

The PT comprised a controlled 2-week introduction phase of body weight exercises and aerobic conditioning. In Weeks 3 and 4, the intensity of PT was increased, and Weeks 5 and 6 then focused on functional fitness and conditioning, including increased load carriage with field packs and weapons. A total of 18, 90-min sessions,

including warm-up and cool-down, were allocated to PT over the 6-week period, and included a combination of aerobic interval running, strength training, circuits, swimming, and bike–boxing–rowing intervals (Table 2). The full detail of the JOIC training programme has been described previously (Edgar et al., 2020).

## 2.6 | Fitness testing

The JOIC fitness evaluation was conducted by NZDF Physical Training Instructors before and after the course. This evaluation consisted of three key components: (a) 2.4-km time-trial run; (b) maximum press-ups; and (c) maximum curl-ups. The 2.4-km road run, which has been shown to provide an effective evaluation of aerobic fitness (Booth et al., 2006; Burger et al., 1990), was completed on a sealed flat road. The run was conducted in a similar fashion to that described by Knapik et al. (2006), where all participants started together, but individual effort was assessed by participants completing the distance in the quickest time possible. Run times were measured via stopwatch to the nearest second (Edgar et al., 2020).

Curl-ups, as used by Vera-Garcia et al., (2000), provided an evaluation of local muscular-endurance of the core where repetitions were completed until failure. The curl-up was performed with participants in a supine position with knees bent flexed at 90° and feet flat on the floor. Hands were held in a fist with arms straight. Hands slid up the thigh until the wrist met the apex of the knee. Hands then slid back down the thigh until the shoulder blades and shoulders touched the ground. A repetition was counted by the instructor every time the wrist reached the apex of the knee until failure. There was no time limit on repetitions, but they were completed in

TABLE 2 JOIC physical and military training programme outline

Variation	Activity	Duration	Number of sessions per week						Total
			Wk 1	Wk 2	Wk 3	Wk 4	Wk 5	Wk 6	
<b>PT</b>									
1	Aerobic interval running	90 min	1	1	1	1	1		5
2	Circuit training (strength endurance)	90 min	1	1	1	1	1		5
3	Swimming/Pool circuit	90 min	1	1	1	1		1	5
4	Stretch, mobility & recovery flush	90 min	1	1		1		1	4
<i>Intensity varied between: High/Medium/Low</i>									
<b>Military training</b>									
1	Drill (Parade ground)	30–60 min	3	2	3	2	2	3	15
2	Weapons training	4 hr +	1	4	3	2	2	3	15
3	Land navigation	3–6 hr		1	2	1			4
4	Sea survival	24 hr					1		1
5	Bush craft	6 hr		1	1	1			3
6	Tactical field exercise	5 days						1	1


A 10-min 06:00 hours early-morning activity (EMA) was also conducted daily, including stretching, mobility and cognitive reaction games.

a continuous fashion with a pause of only 1–2 s between attempts (Edgar et al., 2020).

Press-ups were used to assess upper-body muscular-endurance similar to the protocol outlined by Booth et al. (2006) and Knapik et al. (2006), and were performed on a flat wooden gymnasium surface. Hands were placed on a line in the prone press-up position just slightly wider than shoulder width. A “ready” cue was then given where the body position was adjusted to the start position of arms straight, feet shoulder width apart and the head looking downward. From the start position, the body was lowered eccentrically with a straight-line maintained between the shoulders and heels, until the elbows were at 90°. During the concentric phase, arms were extended until straight while maintaining the back and head positions. Repetitions were completed to fatigue in a continuous fashion and counted by the instructor every time the full range of motion was completed. For both the press-ups and curl-ups, one warning was given for an incomplete repetition, prior to participants being stopped (Edgar et al., 2020).

## 2.7 | Statistical analysis

Scores are shown as mean  $\pm$  SD values unless stated otherwise. We calculated the average sleep duration of all 22 participants over the duration of 36 nights and at the mid-point of total sleep, two quantile groups were identified: UNDERS; averaging  $<6:15$  hr:min of sleep per night,  $n = 11$ ; and OVERS; averaging more than  $6:15$  hr:min sleep per night,  $n = 11$ . The initial intention was to split the sleep groups by those obtaining  $> 7$  hr (the recommended sleep duration per night for adults) compared with those obtaining  $<7$  hr per night. However, given all participants obtained  $<7$  hr per night, the decision was made to split the group in half for further analysis from the median of  $6:15$  hr:min. All statistical analyses were performed using the Statistical Package for Social Science (V. 22.0, SPSS), with statistical significance set at  $p \leq .05$ . To examine whether there were any differences between groups, Group (UNDERS and OVERS)  $\times$  Time (pre- and post- or weekly) two-way multivariate analysis of variance (MANOVAs) were performed on the pre- to post-performance data and weekly subjective wellness data (mood, stress, soreness and fatigue). A Bonferroni adjustment for multiple pairwise comparisons was applied if significant main effects were detected. Analysis of the distribution of residuals was verified visually with histograms, and also using the Shapiro–Wilk test of normality. Magnitudes of the standardized effects between groups were calculated using Cohen's  $d$ , and interpreted using thresholds of 0.2, 0.5 and 0.8 for *small*, *moderate* and *large*, respectively (Cohen, 1988). Effects were deemed *unclear* if the 95% confidence intervals overlapped the *small* thresholds for both positive and negative effects ( $d \pm 0.2$ ). Correlation coefficients (Pearson's  $r$ -values) were determined for the whole group ( $n = 22$ ) to describe associations between sleep and physical performance, and interpreted using thresholds of .00–.19 *very weak*, .20–.39 *weak*, .40–.59 *moderate*, .60–.79 *strong*, and .80–1.0 *very strong* (Evans, 1996).



**FIGURE 1** Percentage performance improvement by group over the 6-week Joint Officer Induction Course (JOIC). Black bars: OVERS ( $> 6:15$  hr:min per night) and white bars UNDERS ( $<6:15$  hr:min). S, small difference between groups. Error bars are 95% confidence intervals

## 3 | RESULTS

The participants were a randomly selected, representative sample from the overall cohort size of  $n = 94$  (Table 1). We were unable to test all 94 participants due to equipment and personnel constraints. From all ( $n = 22$ ) who started the January/February 2019 JOIC, no trainees withdrew from the course or the study. All participant sleep data from every day were included, no data were missed for weekly wellbeing monitoring, and all participants completed both pre- and post-physical performance testing. The average sleep duration across the entire group ( $n = 22$ ) for the JOIC 6-week training course was  $6:10 \pm 0:28$  hr:min. Total weekly sleep reduced by  $6 \pm 2\%$  ( $p = .01$ ) from Week 1 to 6, the equivalent of a  $22 \pm 10$  min per night reduction in total sleep duration. The OVERS group slept on average  $6:27 \pm 9.0$  hr:min, compared with the UNDERS group averaging  $5:51 \pm 28.5$  hr:min, with the OVERS group accumulating  $22:20$  hr:min more sleep than the UNDERS group over the 6-week period ( $p < .01$ ).

The MANOVA detected a significant time effect across all trainees pre-post, regardless of group, for 2.4-km run time, press-ups and curl-ups (all  $p \leq .01$ ), but no significant group  $\times$  time interactions were detected ( $p > .05$ ). However, effect size analysis identified an overall performance improvement favouring OVERS in the 2.4-km run ( $d = 0.29$ , *small*) and press-ups ( $d = -0.30$ , *small*), with a *trivial* difference in the curl-ups ( $d = -0.12$ ). The OVERS group improved 14.9 s more than the UNDERS group (59.8 versus 44.9 s) in the 2.4-km run, and by 1.5 more repetitions (4.7 versus 3.2 reps) in the press-ups (Figure 1; Table 3).

Subjective well-being data for the 6-week training period demonstrated significant and *large* effects in favour of the OVERS group for fatigue and soreness at Weeks 1, 3 and 4. While not

TABLE 3 Mean  $\pm$  SD pre- to post-performance test values for UNDERS (<6 hr 15 min sleep per night,  $n = 11$ ) and OVERS (> 6 hr 15 min sleep per night,  $n = 11$ ), and group  $\times$  time interactions ( $p$ -value and effect size)

	2.4-km run time (s)				Press-ups (repetitions)				Curl-ups (repetitions)			
	Pre-	Post-	$p$ -value	Effect size (d)	Pre-	Post-	$p$ -value	Effect size (d)	Pre-	Post-	$p$ -value	Effect size (d)
UNDERS	601 $\pm$ 75	556 $\pm$ 39	<.01		29 $\pm$ 6	32 $\pm$ 8	<.01		40 $\pm$ 16	44 $\pm$ 17	.05	
OVERS	629 $\pm$ 47	569 $\pm$ 42	<.01		27 $\pm$ 8	31 $\pm$ 8	<.01		39 $\pm$ 22	43 $\pm$ 19	.09	
Group $\times$ time interaction			.37	0.29, Small			.34	-0.30, Small			.79	-0.12, Trivial

statistically significant, there were also *moderate* effects favouring the OVERS compared with UNDERS for stress and mood in Weeks 3 and 4 (Figure 2).

The correlation between sleep and performance metrics and sleep and wellbeing measures was investigated. Regarding performance variables, *moderate* relationships were observed between time in bed and faster 2.4-km run time ( $r = -.47$ ); for sleep-onset latency and press-ups ( $r = .48$ ), and for wake after sleep onset and press-ups ( $r = -.47$ ). Only *weak* or *very weak* relationships were found for all other measures.


## 4 | DISCUSSION

The main results from this study showed that sleeping on average more than 6:15 hr:min per night lead to *small* gains in physical performance measures and had beneficial effects on aspects of subjective well-being in officer trainees over a 6-week training period when compared with sleeping less than 6:15 hr:min per night. The study reinforces previous research, and demonstrates that the recommended 7-9 hr of sleep generally does not occur in the international military personnel (Good et al., 2020; Lentino et al., 2013; Moore et al., 2020). Longer sleep duration in the current study correlated strongly to faster 2.4-km run times, improved number of press-ups, and a shift toward positive fatigue, mood, soreness and stress measures. These data indicate that more time in bed will likely support physical performance adaptation and ongoing physical development in individuals with sleep durations below the recommendation of 7-9 hr (Watson et al., 2015).

Interestingly, the performance improvement observed in the current study was gained in a state similar to that previously described as "short sleep duration", where military members consistently sleep on or around 6 hr per night (Good et al., 2020). While there were no significant differences observed between groups for performance measures, there were *small* effect sizes in favour of the OVERS group for 2.4-km run and press-ups.

Previous military research by Ritland (2019) found a positive relationship between sleep extension (1.36  $\pm$  0.7 hr) and performance benefits in psychomotor vigilance, executive functioning, standing broad jump distance, and motivation levels in officers under training. In a physically demanding professional rugby environment, a ~1-hr sleep extension (from 6:52 to 7:35 hr:min) also showed improved reaction times in a 5-min psychomotor vigilance response test (Swinbourne et al., 2018). Although sleep architecture was not measured in the current study, the aerobic improvement seen in the OVERS group in the current study could potentially be related to an increase of growth hormone release and its relationship to physiological recovery (Dattilo et al., 2011). Growth hormone levels have been shown to effect physical performance, aerobic capacity and, specifically,  $VO_2$  max; thus, it is plausible that these physiological processes have supported muscle recovery and growth, and the observed increase in press-ups and run times (O'Donnell et al., 2018; Widdowson et al., 2009).

Subjective wellbeing in the current study was affected by the quantity of sleep acquired between groups. The reductions in



**FIGURE 2** Weekly subjective wellbeing monitoring of Fatigue, Stress, Soreness and Mood by group over the 6-week Joint Officer Induction Course (JOIC). Solid line; OVERS (> 6:15 hr:min per night) and dashed line; UNDERS (<6:15 hr: min). \*Significant difference between groups ( $p < .05$ ); L, large effect; M, moderate effect. Error bars represent standard deviations

perceived fatigue and soreness in favour of the OVERS group in Weeks 3 and 4 were recorded during stages of high physical and cognitive demand. These observations highlight the important relationship between sleep and enhanced physical and mental wellbeing (Charest & Grandner, 2020; Lentino et al., 2013). Positive outcomes on Stress and Mood scores were also seen in the OVERS group, with reduced stress in Week 4 corresponding with lower soreness, and improved Mood in Week 1. Our data support previous research from Good et al. (2020) that also outlined the strong association between sleep quality and perception of stress and fatigue in military populations, and the association between sleep quantity, cognitive function and reduced physical capacity. In a similar operational environment, McGillis et al. (2017) found that wildland attack firefighters physical performance could be maintained in the initial stages of extended periods of poor sleep and broken shifts (~5 hr a day over 14–18 days), but as days progressed, poor judgement, deflated mood, increased fatigue, muscle soreness, and a decrease in reaction time and physical performance did occur.

Similar to the findings of Moore et al. (2020), the current study observed short sleep duration in this initial stage of military training due to non-standard shift work schedules and routinely participating in demanding and highly variable daily schedules of > 12 hr in duration across several days. A study by Miller and Shattuck (2005), with similar aged officer trainees and sleep to the current study, found that US Military Academy Cadets sleep ranged from 5 hr 50 min to 6 hr 32 min during initial training, and trainees often struggled with intense PT. It has also been acknowledged that

military training, even at academies and officer training schools similar to the JOIC environment, is characterized by highly demanding physical and academic training loads with limited sleep opportunities (Moore et al., 2020). Further to this finding, research conducted by the US Naval postgraduate school over the last decade highlights a common trend of soldiers, sailors and marines worldwide accumulating high levels of sleep debt (Miller et al., 2012), supported by the results of the current study.

Within the constraints of conducting research in the military environment, we acknowledge a limitation in the current study was that the time available to sleep was confined to a specific window from “lights out” at 2,200 hr to “wake-up” at 0,545 hr. This narrow window of sleep opportunity led to a relatively minor difference in sleep duration between the OVERS and UNDERS groups. It is possible that if the sleep and wake times were not set, self-selected differences in sleep duration might have been more pronounced and had a greater impact on the differences in the outcome variables. Future research in military settings should consider comparing groups where different sleep opportunities are set (e.g. <6 hr versus the recommended 7–9 hr) via manipulating sleep and wake times, while measuring physical, cognitive and wellbeing variables.

In conclusion, results from this study have demonstrated that in two groups of trainees who were grouped by sleep duration derived from wrist-actigraphy, a non-significant but *small* improvement in aerobic fitness and press-up performance was seen in recruits sleeping more than 6:15 hr:min compared with those sleeping <6:15 hr:min per night. The group sleeping approximately 37 min

more per night, and thus 22:20 hr:min over the duration of the training course, also showed benefits in aspects of perceived well-being (fatigue, soreness, mood and stress). Thus, even a modest increase in sleep duration in a short sleeping cohort may result in enhanced physical performance and perceived wellbeing.

## ACKNOWLEDGEMENTS

The authors wish to acknowledge the support and assistance of the New Zealand Defence Force, Joint Service Component Commander GRPCAPT P. Franken, Woodbourne Base Commander WGCDCR B. Pothan, and JOIC Chief Instructor SQDLDR B. Warren. Special thanks also to the JOIC training staff and all officer trainees who participated in this study.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest associated with this work.

## AUTHOR CONTRIBUTIONS

DE designed the study, collected the data, analysed the data and wrote the final version of the manuscript. NG designed the study and contributed to the writing of the manuscript. CMB & MD designed the study, analysed the data and contributed to the writing of the manuscript. JZ contributed to the data collection and analysis, and writing of the manuscript.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## ORCID

David T. Edgar  <https://orcid.org/0000-0003-4829-803X>

Christopher Martyn Beaven  <https://orcid.org/0000-0003-2900-7460>

Jennifer L. Zaslona  <https://orcid.org/0000-0002-6954-1150>

Matthew W. Driller  <https://orcid.org/0000-0002-9990-8830>

## REFERENCES

- Booth, C. K., Probert, B., Forbes-Ewan, C., & Coad, R. A. (2006). Australian army recruits in training display symptoms of overtraining. *Military Medicine*, 171(11), 1059–1064. <https://doi.org/10.7205/MILMED.171.11.1059>
- Brown, A., Berry, R., & Schmidt, A. (2013). Sleep and military members: Emerging issues and nonpharmacological intervention. *Sleep Disorders*, 2013, 1–6. <https://doi.org/10.1155/2013/160374>
- Burger, S. C., Bertram, S. R., & Stewart, R. I. (1990). Assessment of the 2.4 km run as a predictor of aerobic capacity. *South African Medical Journal*, 78(9), 327–329.
- Charest, J., & Grandner, M. A. (2020). Sleep and athletic performance: Impacts on physical performance, mental performance, injury risk and recovery, and mental health. *Sleep Medicine Clinics*, 15(1), 41–57. <https://doi.org/10.1016/j.jsmc.2019.11.005>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. Routledge.
- Dattilo, M., Antunes, H., Medeiros, A., Neto, M., Souza, H., Tufik, S., & De Mello, M. (2011). Sleep and muscle recovery: Endocrinological and molecular basis for a new and promising hypothesis. *Medical Hypotheses*, 77(2), 220–222. <https://doi.org/10.1016/j.mehy.2011.04.017>
- Driller, M., O'Donnell, S., & Tavares, F. (2017). What wrist should you wear your actigraphy device on? Analysis of dominant vs. non-dominant wrist actigraphy for measuring sleep in healthy adults. *Sleep Science*, 10(3), 132–135.
- Edgar, C. D. T., Gill, N. D., & Driller, M. W. (2020). Physical characteristics of New Zealand army, navy and airforce officer trainees' over a 6-week joint officer induction course. *The Journal of Sport and Exercise Science*, 4(2), 63–71.
- Evans, J. (1996). *Straightforward statistics for the behavioral sciences*. Thomson Brooks/Cole Publishing Co.
- Good, C. H., Brager, A. J., Capaldi, V. F., & Mysliwiec, V. (2020). Sleep in the United States military. *Journal of Neuropsychopharmacology*, 45(1), 176–191. <https://doi.org/10.1038/s41386-019-0431-7>
- Hooper, S., & Mackinnon, L. (1995). Monitoring overtraining in athletes. *Sports Medicine*, 20(5), 321–327. <https://doi.org/10.2165/00007256-199520050-00003>
- Knapik, J., Darakjy, S., Hauret, K., Canada, S., Scott, S., Rieger, W., Marin, R., & Jones, B. (2006). Increasing the physical fitness of low-fit recruits before basic combat training: An evaluation of fitness, injuries, and training outcomes. *Military Medicine*, 171(1), 45–54. <https://doi.org/10.7205/MILMED.171.1.45>
- Kushida, C. A., Chang, A., Gadkary, C., Guilleminault, C., Carrillo, O., & Dement, W. C. (2001). Comparison of actigraphic, polysomnographic, and subjective assessment of sleep parameters in sleep-disordered patients. *Sleep Medicine*, 2(5), 389–396.
- Lentino, C., Purvis, D., Murphy, K., & Deuster, P. (2013). Sleep as a component of the performance triad: The importance of sleep in a military population. *US Army Medical Department Journal*, 98–107.
- Lovalekar, M., Sharp, M. A., Billing, D. C., Drain, J. R., Nindl, B. C., & Zambraski, E. J. (2018). International consensus on military research priorities and gaps-survey results from the 4th International Congress on Soldiers' Physical Performance. *Journal of Science and Medicine in Sport*, 21, 1125–1130. <https://doi.org/10.1016/j.jsams.2018.05.028>
- McGillis, Z., Dorman, S. C., Robertson, A., Larivière, M., Leduc, C., Eger, T., Oddson, B. E., & Larivière, C. (2017). Sleep quantity and quality of Ontario wildland firefighters across a low-hazard fire season. *Journal of Occupational Environmental Medicine*, 59(12), 1188. <https://doi.org/10.1097/JOM.0000000000001175>
- McHugh, M. L. (2012). Interrater reliability: The kappa statistic. *Journal of Biochemistry Medica*, 22(3), 276–282. <https://doi.org/10.11613/BM.2012.031>
- Miller, N. L., Matsangas, P., & Kenney, A. (2012). *The role of sleep in the military: Implications for training and operational effectiveness*. Naval Postgraduate School Monterey CA Dept of Operations Research.
- Miller, N. L., & Shattuck, L. G. (2005). Sleep patterns of young men and women enrolled at the United States Military Academy: Results from year 1 of a 4-year longitudinal study. *Sleep*, 28(7), 837–841. <https://doi.org/10.1093/sleep/28.7.837>
- Miller, N. L., Shattuck, L. G., & Matsangas, P. (2010). Longitudinal study of sleep patterns of United States Military Academy cadets. *Sleep*, 33(12), 1623–1631.
- Miller, N. L., Shattuck, L. G., & Matsangas, P. (2011). Sleep and fatigue issues in continuous operations: A survey of US army officers. *Behavioral Sleep Medicine*, 9(1), 53–65. <https://doi.org/10.1080/15402002.2011.533994>
- Moore, B., Brock, M., Brager, A., Collen, J., LoPresti, M., & Mysliwiec, V. (2020). Posttraumatic stress disorder, traumatic brain Injury, sleep, and performance in military personnel. *Journal of Sleep Medicine Clinics*, 15, 87–100. <https://doi.org/10.1016/j.jsmc.2019.11.004>
- Morgenthaler, T., Alessi, C., Friedman, L., Owens, J., Kapur, V., Boehlecke, B., Brown, T., Chesson, A., Coleman, J., Lee-Chiong, T., Pancer, J., & Swick, T. J. (2007). Practice parameters for the use of actigraphy in the assessment of sleep and sleep disorders: An update for 2007. *Journal of Sleep*, 30(4), 519–529. <https://doi.org/10.1093/sleep/30.4.519>

- O'Donnell, S., Beaven, C. M., & Driller, M. W. (2018). From pillow to podium: A review on understanding sleep for elite athletes. *Nature & Science of Sleep*, *10*, 243–253.
- Owens, J. (2014). Insufficient sleep in adolescents and young adults: An update on causes and consequences. *Pediatrics*, *134*(3), e921–e932. <https://doi.org/10.1542/peds.2014-1696>
- Quante, M., Kaplan, E., Cailler, M., Rueschman, M., Wang, R., Weng, J., Taveras, E., & Redline, S. (2018). Actigraphy-based sleep estimation in adolescents and adults: A comparison with polysomnography using two scoring algorithms. *Nature and Science of Sleep*, *10*, 13–20.
- Ritland, B., Simonelli, G., Gentili, R., Smith, J., He, X., Mantua, J., Balkin, T., & Hatfield, B. (2019). Effects of sleep extension on cognitive/motor performance and motivation in military tactical athletes. *Sleep Medicine*, *58*, 48–55. <https://doi.org/10.1016/j.sleep.2019.03.013>
- Swinbourne, R., Miller, J., Smart, D., Dulson, D., & Gill, N. (2018). The effects of sleep extension on sleep, performance, immunity and physical stress in rugby players. *Sports*, *6*(2), 42. <https://doi.org/10.3390/sports6020042>
- Tryon, W. (2008). Methods of measuring human activity. *Journal of Behavior Analysis in Health, Sports, Fitness*, *1*(1), 58. <https://doi.org/10.1037/h0100369>
- Vera-Garcia, F. J., Grenier, S. G., & McGill, S. M. (2000). Abdominal muscle response during curl-ups on both stable and labile surfaces. *Physical Therapy*, *80*(6), 564–569. <https://doi.org/10.1093/ptj/80.6.564>
- Watson, N. F., Badr, M. S., Belenky, G., Bliwise, D. L., Buxton, O. M., Buysse, D., Dinges, D. F., Gangwisch, J., Grandner, M. A., Kushida, C., Malhotra, R. K., Martin, J. L., Patel, S. R., Quan, S. F., Tasali, E., Twery, M., Croft, J. B., Maher, E., Barrett, J. A., ... Heald, J. L. (2015). Joint consensus statement of the American Academy of sleep medicine and sleep research society on the recommended amount of sleep for a healthy adult: Methodology and discussion. *Sleep*, *38*(8), 1161–1183. <https://doi.org/10.5665/sleep.4886>
- Widdowson, W. M., Healy, M. L., Sonksen, P. H., & Gibney, J. (2009). The physiology of growth hormone and sport. *Growth Hormone & IGF Research*, *19*(4), 308–319. <https://doi.org/10.1016/j.ghir.2009.04.023>
- Williams, S., Collen, J., Wickwire, E., Lettieri, C., & Mysliwiec, V. (2014). The impact of sleep on soldier performance. *Current Psychiatry Reports*, *16*(8), 458–471. <https://doi.org/10.1007/s11920-014-0459-7>

**How to cite this article:** Edgar DT, Gill ND, Beaven CM, Zaslona JL, Driller MW. Sleep duration and physical performance during a 6-week military training course. *J Sleep Res*. 2021;00:e13393. <https://doi.org/10.1111/jsr.13393>